

Random Walk and Diffusion on a Smash Line Algebra

Demosthenes Ellinas^{*} and Ioannis Tsohantjis^{**}

Department of Sciences
Technical University of Crete
GR-73 100 Chania Crete Greece

Abstract

Working withing the framework of Hopf algebras, a random walk and the associated diffusion equation are constructed on a space that is algebraically described as the merging of the real line algebra with the anyonic line algebra. Technically this merged structure is a smash algebra, namely an algebra resulting by a braided tensoring of real with anyonic line algebras. The motivation of introducing the smashing results from the necessity of having non commuting increments in the random walk. Based on the observable-state duality provided by the underlying Hopf structure, the construction is cast into two dual forms: one using functionals determined by density probability functions and the other using the associated Markov transition operator. The ensuing diffusion equation is shown to possess triangular matrix realization. The study is completed by the incorporation of Hamiltonian dynamics in the above random walk model, and by the construction of the dynamical equation obeyed by statistical moments of the problem for generic entangled density functions.

^{*} Email: ellinas@science.tuc.gr

^{**} Email: ioannis2@otenet.gr

1 Introduction

In recent years considerable amount of work has been devoted to generalization of concepts of random walks and diffusions to spaces described by algebras of functions or operators, see e.g. the general reviews [24],[22],[19],[13]. The mathematical framework mostly adopted for such generalizations is that of Hopf algebras of various kinds and their braided versions ([1], [28]). Remarkably some of the basic features of random walks and diffusion processes, such as Markov property, convolution, statistical (in)dependence, diffusion as limit of random walk [20], solution of diffusion equations by classes of moment polynomials ([12], [8]) etc, have their analogue in the generalized framework. Especially the notion of statistical dependence of steps of a random walk, expressed as the commutation property of increments of walk and the analogue of the central limit theorem have been extended from the commuting case [16], to anticommuting case of fermionic increments [30], to the q -deformed case ([26], [18]), to the case of free products ([29], [27]) as well as to case of superalgebras [2]. A general approach to independence based on the coproduct of coalgebras [25], can be used to study random walks on braided structures ([21], [15], [9]).

Braiding or smashing defined on algebras, coalgebras, bialgebras and Hopf algebras, has recently been systematized and a unification of the various braiding notions has been given ([4], [3]). This compact formulation of smashing products has been used here to introduce the so called smash line algebra $\Omega = A \otimes B$, and to study non commutative random walk and the associated diffusion equation on it. In this present work we undertake a full investigation of the random walk defined on the space ensuing from the merging of the real line A with the anyonic line B [21]. Endowing the algebra of functions of this composite space Ω with a smash product, results into smash line algebra. Due to smashing/braiding, the increments of the walk along the subspace of real line A are commuting, the increments along the subspace of the anyonic line B are q -commuting, while for random steps in the total space Ω the increments are Q -commuting. The outline of the paper has as follows.

In section one, we start with the tensor product algebra of the real line and the anyonic line algebra and proceed to establish a mathematically consistent merging of these two in order to form a braided algebra i.e. a smash algebra. This tensor product algebra is also shown to possess a bialgebra structure whose n -fold coproduct will provide us with the notion of the n -th

step of random walk we shall consider. In closing the section we explain the need of introducing the braiding in that it leads, as it is shown, to a type of non commutativity among the steps of the random walk. In section two the diffusion limit of the random walk on the smash line algebra is obtained and the associated diffusion equation is studied as well as the moment generating function of the random variable of the walk. Section three is devoted to the non stationary random walk on the smash line algebra that expresses the dynamical coupling of the random motion with a general Hamiltonian dynamics. Towards a physical interpretation of the constructed random motion on the variables of the smash line algebra, section four provides a matrix representation of these variables and cast the diffusion equation in the form of a coupled spin-oscillator type of equation of motion, for which the solution is obtained in some special cases. Section five utilizing the state-observable duality built in the Hopf algebra structure of the random walk and gives a brief construction of the transition operator of the walk seen it as a markov process. In next section six, we study the effect of the entanglement i.e statistical correlations between the partial walks of the x variable and the ξ variable, expressed as the non factorizability of the probability density function ruling the random walk of the smash line algebra, and investigate, in the terms of the canonical and q -deformed (for q root of unity) Heisenberg algebra, the structure of the ensuing equations of motion for the statistical moments in the diffusion limit. The last section summarized the results and concludes the paper.

2 The smash line algebra

Consider two real associative algebras $\mathcal{A}$ and $\mathcal{B}$ such that $\mathcal{A} = \mathbf{R}[[x]]$ is the real line algebra, consisting of formal power series in one variable generated by the element x with $\{1, x, x^2, \dots\}$ as linear basis, and $\mathcal{B} = \mathbf{R}[\xi]/\xi^N$, the anyonic line algebra consisting of polynomials in one variable of degree $N-1$, generated by the element ξ with basis $\{1, \xi, \xi^2, \dots, \xi^{N-1}\}$ and $\xi^N = 0$. Both these algebras are equipped with a Hopf algebra and a smash Hopf algebra structure which will be examined in turn.

The algebra $\mathcal{A} = \mathcal{A}(\mu_{\mathcal{A}}, \Delta_{\mathcal{A}}, u_{\mathcal{A}}, \epsilon_{\mathcal{A}})$ is a commuting, cocommuting, and coassociative algebra with product, unit, coproduct, and counit respectively

defined as

$$\begin{aligned}
\mu_{\mathcal{A}}(x \otimes y) &= xy, \\
u_{\mathcal{A}}(c) &= cI_{\mathcal{A}}, \\
\Delta_{\mathcal{A}}(x) &= x \otimes I_{\mathcal{A}} + I_{\mathcal{A}} \otimes x, \\
\varepsilon_{\mathcal{A}}(x) &= 0, \text{ and } \varepsilon_{\mathcal{A}}(I_{\mathcal{A}}) = 1
\end{aligned} \tag{1}$$

for all x, y in $\mathcal{A}$, $c \in \mathbf{R}$ and $I_{\mathcal{A}}$, being the identity element. Considering the trivial braiding or twist map τ on $\mathcal{A}$ such that $\tau(x \otimes y) = y \otimes x$, $\forall x, y \in \mathcal{A}$, we can introduce a trivial smash Hopf algebra structure on $\mathcal{A} \otimes \mathcal{A}$, denoted hereafter as $\mathcal{A}' \equiv \mathcal{A}_{\tau} \bowtie_{\tau} \mathcal{A}$, [3, 4] with smash product, unit, smash co-product and co-unit given respectively by

$$\begin{aligned}
\mu_{\mathcal{A}'} &= (\mu_{\mathcal{A}} \otimes \mu_{\mathcal{A}}) \circ (id_{\mathcal{A}} \otimes \tau \otimes id_{\mathcal{A}}), \\
u_{\mathcal{A}'}(1) &= I_{\mathcal{A}'} = I_{\mathcal{A}} \otimes I_{\mathcal{A}} \\
\Delta_{\mathcal{A}'} &= (id_{\mathcal{A}} \otimes \tau \otimes id_{\mathcal{A}}) \circ (\Delta_{\mathcal{A}} \otimes \Delta_{\mathcal{A}}) \\
\varepsilon_{\mathcal{A}'} &= \varepsilon_{\mathcal{A}} \otimes \varepsilon_{\mathcal{A}}.
\end{aligned} \tag{2}$$

The smash product $\mu_{\mathcal{A}'}$ can easily be extended to an associative product on the n -fold tensor product $\mathcal{A}^{\otimes n} \equiv \mathcal{A}^n = \mathcal{A} \otimes \mathcal{A} \otimes \dots \otimes \mathcal{A}$, ($\mathcal{A}$ being taken n times) given by

$$\mu_{\mathcal{A}^n} = (\mu_{\mathcal{A}^{n-1}} \otimes \mu_{\mathcal{A}}) \circ_{k=1}^{2n-3} (id_{\mathcal{A}}^{\otimes 2n-2-k} \otimes \tau \otimes id_{\mathcal{A}}^{\otimes k}).$$

Then, with $x_i = I_{\mathcal{A}} \otimes \dots \otimes x \otimes \dots \otimes I_{\mathcal{A}}$, where x is in the i -th position in $\mathcal{A}^n$ can be verified that the braiding relations

$$x_i x_j = x_j x_i \text{ for all } i, j$$

are satisfied.

The algebra $\mathcal{B} = \mathcal{B}(\mu_{\mathcal{B}}, u_{\mathcal{B}}, \Delta_{\mathcal{B}}, \varepsilon_{\mathcal{B}})$ is an N -potent, commuting, cocommuting and coassociative algebra with product, coproduct, unit and counit respectively defined as

$$\begin{aligned}
\mu_{\mathcal{B}}(\xi \otimes \eta) &= \xi \eta, \\
u_{\mathcal{B}}(c) &= cI_{\mathcal{B}}, \\
\Delta_{\mathcal{B}}(\xi) &= \xi \otimes I_{\mathcal{B}} + I_{\mathcal{B}} \otimes \xi, \\
\varepsilon_{\mathcal{B}}(\xi) &= 0, \text{ and } \varepsilon_{\mathcal{B}}(I_{\mathcal{B}}) = 1
\end{aligned} \tag{3}$$

for all ξ, η in $\mathcal{B}$, $c \in \mathbf{R}$ and $I_{\mathcal{B}}$, being the identity element. Let us consider this time the braiding or twist map ψ on $\mathcal{B}$ such that

$$\psi(\xi \otimes \eta) = q\eta \otimes \xi, \quad \forall \xi, \eta \in \mathcal{B} \quad (4)$$

where $q = e^{2\pi i/N}$, a N -th root of unity. We can introduce a smash Hopf algebra structure on $\mathcal{B} \otimes \mathcal{B}$, denoted hereafter as $\mathcal{B}' \equiv \mathcal{B}_{\psi} \bowtie_{\psi} \mathcal{B}$, [3, 4] with smash product, unit, smash co-product and co-unit given respectively by

$$\begin{aligned} \mu_{\mathcal{B}'} &= (\mu_{\mathcal{B}} \otimes \mu_{\mathcal{B}}) \circ (id_{\mathcal{B}} \otimes \psi \otimes id_{\mathcal{B}}), \\ u_{\mathcal{B}'}(1) &= I_{\mathcal{B}'} = I_{\mathcal{B}} \otimes I_{\mathcal{B}}, \\ \Delta_{\mathcal{B}'} &= (id_{\mathcal{B}} \otimes \psi \otimes id_{\mathcal{B}}) \circ (\Delta_{\mathcal{B}} \otimes \Delta_{\mathcal{B}}), \\ \varepsilon_{\mathcal{B}'} &= \varepsilon_{\mathcal{B}} \otimes \varepsilon_{\mathcal{B}}. \end{aligned} \quad (5)$$

The smash product $\mu_{\mathcal{B}'}$ can be extended to an associative product on the n -fold tensor product $\mathcal{B}^{\otimes n} \equiv \mathcal{B}^n = \mathcal{B} \otimes \mathcal{B} \otimes \dots \otimes \mathcal{B}$, ($\mathcal{B}$ being taken n times) given by

$$\mu_{\mathcal{B}^n} = (\mu_{\mathcal{B}^{n-1}} \otimes \mu_{\mathcal{B}}) \circ_{k=1}^{2n-3} \circ (id_{\mathcal{B}}^{\otimes 2n-2-k} \otimes \psi \otimes id_{\mathcal{B}}^k).$$

Then, with $\xi_i = I_{\mathcal{B}} \otimes \dots \otimes \xi \otimes \dots \otimes I_{\mathcal{B}}$, where ξ is in the i -th position in $\mathcal{B}^n$ can be verified that the braiding relations

$$\xi_i \xi_j = q \xi_j \xi_i \text{ for } i > j. \quad (6)$$

are satisfied as in [21].

The merging of $\mathcal{A}$ and $\mathcal{B}$ as $\Omega \equiv \mathcal{A} \otimes \mathcal{B}$, where x is embedded as $x \otimes I_{\mathcal{A}}$ and ξ as $I_{\mathcal{B}} \otimes \xi$, admits a bialgebra structure with product, unit, co-product, co-unit respectively given by

$$\begin{aligned} \mu_{\Omega} &= (\mu_{\mathcal{A}} \otimes \mu_{\mathcal{B}}) \circ (id_{\mathcal{A}} \otimes \tau \otimes id_{\mathcal{B}}), \\ u_{\Omega}(1) &= I_{\Omega} = I_{\mathcal{A}} \otimes I_{\mathcal{B}}, \\ \Delta_{\Omega} &\equiv (id_{\mathcal{A}} \otimes \tau \otimes id_{\mathcal{B}}) \circ (\Delta_{\mathcal{A}} \otimes \Delta_{\mathcal{B}}), \\ \varepsilon_{\Omega} &\equiv \varepsilon_{\mathcal{A}} \otimes \varepsilon_{\mathcal{B}}. \end{aligned} \quad (7)$$

By means of the previous relations and the braiding map $\Psi : \Omega \otimes \Omega \rightarrow \Omega \otimes \Omega$ defined by

$$\Psi(x^\alpha \otimes \xi^m \otimes x^\beta \otimes \xi^n) = q^{mn} Q^{\alpha n + \beta m} (x^\beta \otimes \xi^n \otimes x^\alpha \otimes \xi^m), \quad (8)$$

for all $x, y \in \mathcal{A}$ and all $\xi, \eta \in \mathcal{B}$, $Q \in \mathbf{R}$ and $q = e^{2\pi i/N}$, we introduce a smash algebra hereafter denoted by $\Omega' = \Omega \#_\Psi \Omega$. Then Ψ can also be written as

$$\Psi = (id_{\mathcal{A}} \otimes \Psi_{\mathcal{AB}} \otimes id_{\mathcal{B}}) \circ (\Psi_{\mathcal{A}} \otimes \Psi_{\mathcal{B}}) \circ (id_{\mathcal{A}} \otimes \Psi_{\mathcal{AB}} \otimes id_{\mathcal{B}}),$$

where $id_{\mathcal{A}} \otimes \Psi_{\mathcal{AB}} \otimes id_{\mathcal{B}}$ is realized as

$$(id_{\mathcal{A}} \otimes \Psi_{\mathcal{AB}} \otimes id_{\mathcal{B}})(x^\alpha \otimes \xi^m \otimes x^\beta \otimes \xi^n) = Q^{\beta m} x^\alpha \otimes x^\beta \otimes \xi^m \otimes \xi^n$$

and $\Psi_{\mathcal{A}}$, $\Psi_{\mathcal{B}}$ are realized as

$$\begin{aligned} \Psi_{\mathcal{A}}(x^\alpha \otimes x^\beta) &= x^\beta \otimes x^\alpha, \\ \Psi_{\mathcal{B}}(\xi^m \otimes \xi^n) &= q^{mn} \xi^n \otimes \xi^m. \end{aligned}$$

Fig. Diagramatic display of the braiding map

The smash product and unit, are given respectively by

$$\begin{aligned} \mu_{\Omega'} &\equiv (\mu_{\Omega} \otimes \mu_{\Omega}) \circ (id_{\Omega} \otimes \Psi \otimes id_{\Omega}), \\ u_{\Omega'}(1) &= I_{\Omega'} = I_{\Omega} \otimes I_{\Omega}. \end{aligned} \quad (9)$$

The following lemma [4], which is easily verified, provides with the compatibility between the above braiding map Ψ and the associative product $\mu_{\Omega'}$.

Lemma: Let $\Omega = \mathcal{A} \otimes \mathcal{B}$ and the linear braiding map $\Psi : \Omega \otimes \Omega \rightarrow \Omega \otimes \Omega$ given in eq.(8). The algebra $\Omega' = \Omega \#_{\Psi} \Omega$ is a smash product algebra which implies that the following relations hold:

i) Ψ is normal i.e

$$\begin{aligned}\Psi(x^{\alpha} \otimes \xi^m \otimes I_A \otimes I_B) &= I_A \otimes I_B \otimes x^{\alpha} \otimes \xi^m \\ \Psi(I_A \otimes I_B \otimes x^{\alpha} \otimes \xi^m) &= (x^{\alpha} \otimes \xi^m \otimes I_A \otimes I_B)\end{aligned}$$

ii) Ψ satisfies the following relation:

$$\begin{aligned}\Omega^{\otimes 4} \rightarrow \Omega^{\otimes 2} : (I_{\Omega} \otimes \mu_{\Omega}) \circ (\Psi \otimes I_{\Omega}) \circ (I_{\Omega} \otimes \mu_{\Omega} \otimes I_{\Omega}) \circ (I_{\Omega} \otimes I_{\Omega} \otimes \Psi) \\ = (\mu_{\Omega} \otimes I_{\Omega}) \circ (I_{\Omega} \otimes \Psi) \circ (I_{\Omega} \otimes \mu_{\Omega} \otimes I_{\Omega}) \circ (\Psi \otimes I_{\Omega} \otimes I_{\Omega}).\end{aligned}$$

The non commutativity of the algebra of increments can explicitly be seen if we look at the simplest example of the one-step algebra of increment as follows:

Define the one-step increments ω_{ij} , $i, j = 1, 2$ as

$$\begin{aligned}\omega_{11} &= x \otimes \xi \otimes I_A \otimes I_B, \quad \omega_{12} = x \otimes I_B \otimes I_A \otimes \xi, \\ \omega_{21} &= I_A \otimes \xi \otimes x \otimes I_B, \quad \omega_{22} = I_A \otimes I_B \otimes x \otimes \xi,\end{aligned}$$

then it can be easily checked that the following relations hold:

$$\begin{aligned}\omega_{11} \omega_{12} &= qQ\omega_{12}\omega_{11}, \quad \omega_{21}\omega_{11} = Q\omega_{11}\omega_{21}, \quad \omega_{22}\omega_{11} = qQ^2\omega_{11}\omega_{22}, \\ \omega_{12}\omega_{21} &= q\omega_{21}\omega_{12}, \quad \omega_{21}\omega_{22} = qQ\omega_{22}\omega_{21}, \quad \omega_{22}\omega_{12} = Q\omega_{12}\omega_{22}.\end{aligned}$$

Note that the above choice for Ψ reproduces the braiding of $\mathcal{A}$ and $\mathcal{B}$ algebras. The product μ_{Ω} extends to an associative product on the n -fold tensor product $\Omega^n \equiv \Omega^{\otimes n}$, $n \geq 2$, by means of the relation

$$\mu_{\Omega^n} = (\mu_{\Omega^{n-1}} \otimes \mu_{\Omega}) \circ_{k=1}^{2n-3} \circ (id_{\Omega}^{\otimes 2n-2-k} \otimes \Psi \otimes id_{\Omega}^{\otimes k}), \quad (10)$$

and in this way provides the following braiding relations on Ω^n :

$$x_i x_j = x_j x_i, \quad \forall i, j, \quad \xi_i \xi_j = q \xi_j \xi_i, \quad x_i \xi_j = Q \xi_j x_i, \quad \text{for } i > j, \quad (11)$$

where the indices above indicate the position of the embeddings of x and ξ in the respective spaces (e.g. $\xi_2 \xi_1 = (I_A \otimes I_B \otimes I_A \otimes \xi)(I_A \otimes \xi \otimes I_A \otimes I_B) = q(I_A \otimes \xi \otimes I_A \otimes I_B)(I_A \otimes I_B \otimes I_A \otimes \xi) = q\xi_1 \xi_2$).

Finally using eqs. (7,11), we compute the n -th fold coproduct on $x^k \otimes \xi^l \in \Omega$, $k \in \mathbf{Z}_+$, $l \in \{0, 1, \dots, N-1\}$, which is given by:

$$\begin{aligned} \Delta_\Omega^{n-1}(x^k \otimes \xi^l) = & \sum_{i_1 + \dots + i_n = k} \sum_{j_1 + \dots + j_n = l} \begin{pmatrix} k \\ i_1 & \dots & i_n \end{pmatrix} \begin{bmatrix} l \\ j_1 & \dots & j_n \end{bmatrix}_q \\ & \times x^{i_1} \otimes \xi^{j_1} \otimes \dots \otimes x^{i_n} \otimes \xi^{j_n} \end{aligned} \quad (12)$$

where $\Delta_\Omega^m = (\Delta_\Omega \otimes id_\Omega^{\otimes m-1}) \circ \Delta_\Omega^{m-1} = (id^{\otimes m-1} \otimes \Delta_\Omega) \circ \Delta_\Omega^{m-1}$, with $\begin{pmatrix} k \\ i_1 & \dots & i_n \end{pmatrix} = \frac{k!}{i_1! \dots i_n!}$, while the q -binomial coefficient is defined as $\begin{bmatrix} l \\ j_1 & \dots & j_n \end{bmatrix}_q = \frac{[l]_q!}{[j_1]_q! \dots [j_n]_q!}$, where $[l]_q = \frac{q^l - 1}{q - 1}$, $[l]_q! = [1]_q [2]_q \dots [l]_q$.

In the next chapter we will construct a stationary random walk on Ω and derive the diffusion limit equation of motion.

3 Diffusion equation on the smash line algebra

Let $\phi_\Omega : \Omega \longrightarrow \mathbf{C}$, be a linear normalized ($\phi_\Omega(\mathbf{1}_\Omega) = 1$) and positive semi-definite functional from Ω to $\mathbf{C}$, which corresponds to probability density element ρ satisfying the following relations:

$$\phi_\Omega(f) \equiv \langle f \rangle_{\phi_\Omega} = \int \rho f = \langle \phi_\Omega, f \rangle, \quad (13)$$

where $f = f(x, \xi)$ is a general element (observable) of Ω . It is assumed that ϕ_Ω lives in the dual space of Ω , where the product is defined to be the usual convolution operation between probability density functions, by means of the following relations:

$$\phi_\Omega^{\star n}(f) = \langle \phi_\Omega \star \phi_\Omega \star \dots \star \phi_\Omega, f \rangle = \langle \phi_\Omega^{\otimes n}, \Delta^{n-1}(f) \rangle. \quad (14)$$

where $\Delta^{n-1} \equiv \Delta_{\Omega}^{n-1}$ will be used thereafter. If eq.(14) is interpreted as the state of probability function of the random walk after n steps, then we can evaluate the general state after an n -step walk. Consider a general observable element of Ω such as

$$f(x, \xi) = \sum_{l=0}^{N-1} \sum_{k \in \mathbf{Z}_+} c_{kl} x^k \xi^l \equiv \sum_{l=0}^{N-1} c_l(x) \xi^l = \sum_{k \in \mathbf{Z}_+} x^k d_k(\xi) \quad (15)$$

where it should be understood, except if otherwise stated, that $x^k \equiv (x \otimes I_B)^k = (x^k \otimes I_B)$ and $\xi^l \equiv (I_A \otimes \xi)^l = (I_A \otimes \xi^l)$. The last equations indicate that the observable $f(x, \xi)$, can be seen either as an $N - 1$ -degree anyonic polynomial with real analytical function as coefficients ($c_l(x)$), or as a real analytic function with anyonically valued coefficients. In view of the matrix realization of the anyonic variables discussed below, these two alternatives of representing the general observable quantities of the random walk might help us to identify a physical model described by the smash algebra Ω , as a model of interacting classical/boson system to a spin like system. Next using eqs.(10,12,13,14,15), we obtain the convolution

$$\begin{aligned} \phi_{\Omega}^{\star n}(f) = \\ \sum_{k, l \in \mathbf{Z}_+} \sum_{i_1 + \dots + i_n = k} \sum_{j_1 + \dots + j_n = l} \begin{pmatrix} k \\ i_1 & \dots & i_n \end{pmatrix} \begin{bmatrix} l \\ j_1 & \dots & j_n \end{bmatrix}_q c_{kl} \\ \times \phi_{\Omega}(x^{i_1} \otimes \xi^{j_1}) \dots \phi_{\Omega}(x^{i_n} \otimes \xi^{j_n}). \end{aligned} \quad (16)$$

Note that although ϕ_{Ω} is linear functional the convoluted $\phi_{\Omega}^{\star n}$ is a non-linear one. In order to evaluate the above quantity we can assume that

$$\rho_{\Omega}(x, \xi) = \rho_A(x) \otimes \rho_B(\xi). \quad (17)$$

The density matrices $\rho_A(x)$, $\rho_B(\xi)$ can be taken of the form

$$\begin{aligned} \rho_A(x) &= p_1 \delta(x - a) + (1 - p_1) \delta(x + a), \\ \rho_B(\xi) &= p_2 \delta(\xi - \theta) + (1 - p_2) \delta(\xi + \theta), \end{aligned} \quad (18)$$

where as usual p_1 and p_2 are chosen probabilities and the anyonic delta function in $\rho_B(\xi)$ is defined as $\delta(\xi - \theta) = \sum_{i=0}^{N-1} \theta^{N-1-i} \xi^i$ and $\int \delta(\xi + \theta) f(\xi) = f(\theta)$ [21]. This choice allows us to determine explicitly average values of the generating monomials of the algebra after n steps. Note that the value of the size of each random step in Ω is considered in general to be $a \otimes \theta$. For the random walk on the real line in Ω we have that

$$\begin{aligned}
\langle x^m \rangle_{\phi_{\Omega}^{*n}} &= \sum_{s_1 + \dots + s_n = m} \binom{m}{s_1 \dots s_n} \Pi_{l=1}^n \langle x^{s_l} \rangle_{\phi_A} \\
&= \sum_{s_1 + \dots + s_n = m} \binom{m}{s_1 \dots s_n} \Pi_{l=1}^n (p_1 a^{s_l} + (1 - p_1)(-a)^{s_l}) \\
&= \sum_{s_1 + \dots + s_n = m} \binom{m}{s_1 \dots s_n} \Pi_{l=1}^n [p_1 e^{aD_x} \otimes \varepsilon_{\mathcal{B}} + (1 - p_1) e^{-aD_x} \otimes \varepsilon_{\mathcal{B}}]_{|x=0} (x^{s_l} \otimes I_{\mathcal{B}}) \\
&= \sum_{s_1 + \dots + s_n = m} \binom{m}{s_1 \dots s_n} \Pi_{l=1}^n \phi_A(x^{s_l}) \\
&= \sum_{s_1 + \dots + s_n = m} \binom{m}{s_1 \dots s_n} \Pi_{l=1}^n \langle x^{s_l} \rangle_{\phi_A}
\end{aligned} \tag{19}$$

where $D_x = \partial/\partial_x$.

With respect to random walks on the anyonic line in Ω we have that:

$$\begin{aligned}
\langle \xi^t \rangle_{\phi_{\Omega}^{*n}} &= \sum_{r_1 + \dots + r_n = t} \left[\begin{matrix} t \\ r_1 \dots r_n \end{matrix} \right]_q \Pi_{l=1}^n \langle \xi^{r_l} \rangle_{\phi_B} \\
&= \sum_{r_1 + \dots + r_n = t} \left[\begin{matrix} t \\ r_1 \dots r_n \end{matrix} \right]_q \Pi_{l=1}^n (p_2 \theta^{r_l} + (1 - p_2)(-\theta)^{r_l}) \\
&= \sum_{r_1 + \dots + r_n = t} \left[\begin{matrix} t \\ r_l \dots r_n \end{matrix} \right]_q \Pi_{l=1}^n [\varepsilon_{\mathcal{A}} \otimes p_2 e^{\theta D_{\xi}} + (1 - p_2) \varepsilon_{\mathcal{A}} \otimes e^{-\theta D_{\xi}}]_{|\xi=0} (I_{\mathcal{A}} \otimes \xi^{r_l}) \\
&= \sum_{r_1 + \dots + r_n = t} \left[\begin{matrix} t \\ r_l \dots r_n \end{matrix} \right]_q \Pi_{l=1}^n \phi_B(\xi^{r_l}) \\
&= \sum_{r_1 + \dots + r_n = t} \left[\begin{matrix} t \\ r_l \dots r_n \end{matrix} \right]_q \Pi_{l=1}^n \langle \xi^{r_l} \rangle_{\phi_B}
\end{aligned} \tag{20}$$

where the derivative D_{ξ} is defined in [21, 10] as $D_{\xi} f(\xi) = \frac{f(\xi) - f(\xi q)}{(1-q)\xi}$, and satisfies the q -Leibniz rule $D_{\xi}(fg) = (D_{\xi}f)g + (L_q f)D_{\xi}g$, where $L_q f(\xi) = f(q\xi)$.

Finally with respect to simultaneous random walks in both x, ξ directions we obtain using eqs.(19,20):

$$\begin{aligned} & \langle x^k \otimes \xi^l \rangle_{\phi^{\star n}} = \phi^{\star n}(x^k \otimes \xi^l) \\ & = \sum_{i_1+\dots+i_n=k} \sum_{j_1+\dots+j_n=l} \begin{pmatrix} k \\ i_1 & \dots & i_n \end{pmatrix} \begin{bmatrix} l \\ j_1 & \dots & j_n \end{bmatrix}_q \end{aligned} \quad (21)$$

$$\begin{aligned} & \times \prod_{s=1}^n \phi_x(x^{i_s}) \phi_\xi(\xi^{j_s}) \\ & = \phi_x^{\star n}(x^k) \phi_\xi^{\star n}(\xi^l) \end{aligned} \quad (22)$$

where $\phi_x(x^{i_s}) \phi_\xi(\xi^{j_s}) = \langle x^{i_s} \rangle_{\phi_x} \langle \xi^{j_s} \rangle_{\phi_\xi}$.

Let us now compute the system after n steps and its limit as $n \rightarrow \infty$. Using Taylor's expansion, for the form of $\phi_\Omega^{\star n}$ as it can be easily read off from equations (14,19,20), and (21) we have that

$$\begin{aligned} \phi_\Omega^{\star n} &= \phi_A^{\star n} \otimes \phi_B^{\star n} = ([\varepsilon_{\mathcal{A}} + 2a(p_1 - 1/2)D_x + a^2/2!D_x^2 + \dots]^n \\ & \quad \otimes [\varepsilon_{\mathcal{B}} + 2\theta(p_2 - 1/2)D_\xi + \theta^2/[2]_q!D_\xi^2 + \dots]^n) |_{x=0, \xi=0}. \end{aligned} \quad (23)$$

Following [20, 21], we make the following substitutions

$$\begin{aligned} 2a(p_1 - 1/2) &= \frac{c_1 t}{n}, \quad 2\theta(p_2 - 1/2) = \frac{c_2 t}{n}, \\ a^2/2 &= \frac{\alpha_1 t}{n}, \quad \theta^2/[2]_q = \frac{\alpha_2 t}{n}. \end{aligned} \quad (24)$$

Then we take the limit $n \rightarrow \infty$ with $t, c_1, c_2, \alpha_1, \alpha_2$ kept fixed and $t = nd, d$ being the size of the step in time (which in the limit taken $d \rightarrow 0$ as a does), to obtain the continue limit of random walk where the steps are viewed as steps in time. Thus we obtain:

$$\phi_\Omega^\infty(f) = (e^{(c_1 t D_x + \alpha_1 t D_x^2) \otimes \varepsilon_{\mathcal{B}} + \varepsilon_{\mathcal{A}} \otimes (c_2 t D_\xi + \alpha_2 t D_\xi^2)} f) |_{x, \xi=0}, \quad (25)$$

where the limit $(1 + z/n)^n \rightarrow e^z$ for $n \rightarrow \infty$, has been used. The associated density ρ_Ω^∞ can now be obtained by evaluating

$$\phi_\Omega^\infty(\delta) = \int \rho^\infty \delta(x, \xi), \text{ where } \delta(x, \xi) \equiv \delta(x - a) \otimes \delta(\xi - \theta).$$

Then

$$\rho_{\Omega}^{\infty}(a, \theta) = \rho_A^{\infty}(a) \otimes \rho_B^{\infty}(\theta), \quad (26)$$

where

$$\rho_A^{\infty}(a) = (4\pi\alpha_1 t)^{-1} e^{-\frac{(a-c_1 t)^2}{4\alpha_1 t}}, \quad (27)$$

$$\rho_B^{\infty}(\theta) = \sum_{k=0}^{N-1} \theta^{N-1-k} \sum_{l=0}^{l < k/2} \frac{(c_2 t)^{k-l} (\alpha_2/c_2)^l [k]_q!}{l!(k-2l)!}. \quad (28)$$

To obtain the diffusion equation we take a generic f of the form (15) and evaluate:

$$\begin{aligned} \int (\partial_t \rho_{\Omega}^{\infty}) f &= \partial_t \phi_{\Omega}^{\infty}(f) = \phi_{\Omega}^{\infty}((c_1 D_x + \alpha_1 D_x^2 + c_2 D_{\xi} + \alpha_2 D_{\xi}^2) f) \\ &= \int \rho_{\Omega}^{\infty} (c_1 D_x + \alpha_1 D_x^2 + c_2 D_{\xi} + \alpha_2 D_{\xi}^2) f \\ &= \int ((-c_1 D_x + \alpha_1 D_x^2 + c_2 D_{\xi}^* + \alpha_2 D_{\xi}^{*2}) \rho^{\infty}) f, \end{aligned} \quad (29)$$

where we have used the definition of $D_{\xi}^* = -D_{\xi} L_{q^{-1}}$, $L_{q^{-1}}$ being such that $L_{q^{-1}} f(\xi) = f(q^{-1} \xi)$. Then eq.(29) leads to the desired diffusion equation:

$$\partial_t \rho_{\Omega}^{\infty} = ((-c_1 D_x + \alpha_1 D_x^2) \otimes id_B + id_A \otimes (c_2 D_{\xi}^* + \alpha_2 D_{\xi}^{*2})) \rho_{\Omega}^{\infty} \quad (30)$$

whose solution is given in eqs.(26,27,28) .

Before closing this chapter we will quote the generating function of the statistical moments of the two random variables, which are:

$$G(k_1, k_2) \equiv \langle e^{ik_1 x} \otimes e^{ik_2 \xi} \rangle_{\phi} = \sum_{m_1 \geq 0} \sum_{m_2=0}^{N-1} \frac{(ik_1)^{m_1}}{m_1!} \frac{(ik_2)^{m_2}}{m_2!} \langle x^{m_1} \rangle_{\phi} \langle \xi^{m_2} \rangle_{\phi}$$

Then we can obtain the moments as:

$$\begin{aligned} \frac{d^l}{i^l dk_1^l} G(k_1, k_2)|_{k_1=k_2=0} &= \langle x^l \rangle_{\phi}, \\ (1/i) D_{q, k_2} G(k_1, k_2)|_{k_1=k_2=0} &= \langle \xi \rangle_{\phi}, \\ (1/i^l) D_{q, k_2}^l G(k_1, k_2)|_{k_1=k_2=0} &= \langle \xi^l \rangle_{\phi}, \end{aligned} \quad (31)$$

where $D_{q, k_2} f(k_2) = \frac{f(k_2) - f(qk_2)}{(1-q)k_2}$, the q -deformed derivative[10].

4 Non stationary random walk

Consider now the Hamiltonian evolution of a quantity $F = F(x, p, \xi, p_\xi)$, depending in general on the phase space variables x, p and ξ, p_ξ determined by the following Poisson brackets

$$\begin{aligned}\{F, H\} &= \frac{\partial F}{\partial x} \frac{\partial H}{\partial p} - \frac{\partial F}{\partial p} \frac{\partial H}{\partial x} + (-1)^\epsilon \left(\frac{\partial F}{\partial \xi} \frac{\partial H}{\partial p_\xi} + \frac{\partial F}{\partial p_\xi} \frac{\partial H}{\partial \xi} \right) = V_H(F) \\ &= V_H^1(F) + V_H^2(F),\end{aligned}\tag{32}$$

where we have assumed that $\xi^2 = 0$. The parity degree of F , $\deg(F) = \epsilon$ takes values $\epsilon = 0, 1$ to account of the even or odd character of F respectively. The form and action of the operators $V_H^1 = \frac{\partial H}{\partial p} \frac{\partial}{\partial x} - \frac{\partial H}{\partial x} \frac{\partial}{\partial p}$, $V_H^2 = (-1)^\epsilon \left(\frac{\partial}{\partial \xi} \frac{\partial H}{\partial p_\xi} + \frac{\partial}{\partial p_\xi} \frac{\partial H}{\partial \xi} \right)$, can be read off from the above equation. We can thus defined, using t as a time parameter, the action of an evolution operator e^{tV_H} on x_0 , by $e^{tV_H} x_0 = x_t$, and on ξ_0 by $e^{tV_H} \xi_0 = \xi_t$ where x_t and ξ_t are the values of the observables at time t . We shall now proceed to calculate moments after one step walk in Ω , $\phi(x_t^k \otimes \xi_t^l)$, where $k \in \mathbf{Z}_+$, $l = 0, 1$, using the probability densities for time $t = 0$, given as $\rho_0^1(x_0) = p_1 \delta(x_0 - a) + (1 - p_1) \delta(x_0 + a)$, $\rho_0^2(\xi_0) = p_2 \delta(\xi_0 - \theta) + (1 - p_2) \delta(\xi_0 + \theta)$, while the delta functions for the ξ variables are as defined previously. Moreover we shall again assume that $\rho_0(x_0, \xi_0) = \rho_0^1(x_0) \otimes \rho_0^2(\xi_0)$. The reason for these choices will become apparent below. Thus,

$$\begin{aligned}\phi(x_t \otimes \xi_t) &= \langle \rho_0, e^{tV_H^1} x_0 \otimes e^{tV_H^2} \xi_0 \rangle \\ &= \langle e^{-tV_H^1} \rho_0^1, x_0 \rangle \langle e^{-tV_H^2} \rho_0^2, \xi_0 \rangle = \rho^1(x_t) \rho^2(\xi_t) \\ &= \rho_t^1(x_0) \rho_t^2(\xi_0).\end{aligned}\tag{33}$$

Direct evaluation of ρ_t^1 and ρ_t^2 gives

$$\begin{aligned}\rho_t^1(x_0) &= p_1 \delta(x_0 - t \frac{\partial H}{\partial p} - a) + (1 - p_1) \delta(x_0 - t \frac{\partial H}{\partial p} + a) \\ &= p_1 \delta(x_0 - t\lambda - a) + (1 - p_1) \delta(x_0 - t\lambda + a),\end{aligned}\tag{34}$$

$$\rho_t^2(\xi_0) = p_2 \delta(\xi_0 - t \frac{\partial H}{\partial p_\xi} - \theta) + (1 - p_2) \delta(\xi_0 - t \frac{\partial H}{\partial p_\xi} + \theta)\tag{35}$$

$$= p_2 \delta(\xi_0 - t\tilde{\lambda} - \theta) + (1 - p_2) \delta(\xi_0 - t\tilde{\lambda} + \theta).\tag{36}$$

where $\lambda = \frac{\partial H}{\partial p}$ and $\tilde{\lambda} = -\frac{\partial H}{\partial p_\xi}$ and thus, using eqs.(13,33) we finally have

$$\begin{aligned}
\phi(x_t \otimes \xi_t) &= [(p_1 e^{a_t D_{x_0}} \otimes \varepsilon_{\mathcal{B}} + (1 - p_1) e^{a'_t D_{x_0}} \otimes \varepsilon_{\mathcal{B}}) \\
&\quad \times (\varepsilon_{\mathcal{A}} \otimes p_2 e^{\theta_t D_{\xi_0}} + (1 - p_2) \varepsilon_{\mathcal{A}} \otimes e^{\theta'_t D_{\xi_0}})]|_{x_0, \xi_0=0} (x_0 \otimes \xi_0) \\
&= e^{-\lambda t \partial / \partial a} e^{-\tilde{\lambda} t \partial / \partial \theta} \phi_{t=0}(x_0 \otimes \xi_0) \\
&= \phi_t(x_0 \otimes \xi_0)
\end{aligned} \tag{37}$$

where we have set $a_t = a - t\lambda$, $a'_t = -a - t\lambda$, $\theta_t = \theta - t\tilde{\lambda}$ and $\theta'_t = -\theta - t\tilde{\lambda}$. Evaluation of $\phi(f)$ for a generic f of the form given above can be straightforwardly calculated using eq.(37). To derive the continue limit we proceed as before. By Taylor's expansion of ϕ_t and taking $\phi_t^{\star n}$ we have

$$\begin{aligned}
\phi_t^{\star n} &= [\varepsilon_{\Omega} + (2a(p_1 - 1/2) - t\lambda) D_x \otimes \varepsilon_{\mathcal{B}} + \\
&\quad (\frac{a^2}{2} + \frac{(t\lambda)^2}{2} + at\lambda - 2p_1 ta\lambda) D_x^2 \otimes \varepsilon_{\mathcal{B}} + \dots]^n \\
&\quad \times [\varepsilon_{\Omega} + \varepsilon_{\mathcal{A}} \otimes (2\theta(p_2 - 1/2) - t\tilde{\lambda}) D_{\xi} + \\
&\quad \varepsilon_{\mathcal{A}} \otimes (\frac{\theta^2}{2} + \frac{(t\tilde{\lambda})^2}{2} + \theta t\tilde{\lambda} - 2p_2 t\theta\tilde{\lambda}) D_{\xi}^2 + \dots]^n.
\end{aligned} \tag{38}$$

As before we are making the following substitutions

$$\begin{aligned}
2a(p_1 - 1/2) &= \frac{c_1 t}{n}, \quad t\lambda = \lambda \frac{d_1 t}{n}, \quad \frac{a^2}{2} = \frac{\alpha_1 t}{n}, \\
2\theta(p_2 - 1/2) &= \frac{c_2 t}{n}, \quad t\tilde{\lambda} = \tilde{\lambda} \frac{d_2 t}{n}, \quad \frac{\theta^2}{2} = \frac{\alpha_2 t}{n},
\end{aligned} \tag{39}$$

where d_1, d_2 , real constants. Here $D_{\xi}^2 = 0$ and thus the last of eq.(39) is not needed. Taking the limit $n \rightarrow \infty$ as before we obtain for a generic function $f(x, \xi)$

$$\begin{aligned}
\phi^{\infty}(f) &= [e^{((c_1 - \lambda d_1)t D_x + \alpha_1 t D_x^2) \otimes \varepsilon_{\mathcal{B}} + \varepsilon_{\mathcal{A}} \otimes (c_2 - \tilde{\lambda} d_2)t D_{\xi}}]_{|x, \xi=0} f \\
&= (e^{tK} f)_{|x, \xi=0} = \int \rho^{\infty} f = \langle \rho^{\infty}, f \rangle.
\end{aligned} \tag{40}$$

The diffusion equation is immediately implied from the fact that:

$$\langle \partial_t \rho^{\infty}, f \rangle = \int \partial_t \rho^{\infty} f = (e^{tK} K f)_{|x, \xi=0} = \langle \rho^{\infty}, K f \rangle = \langle K^* \rho^{\infty}, f \rangle$$

giving

$$\begin{aligned} \frac{\partial \rho^\infty}{\partial t} = & [-c_1 D_x \otimes id_{\mathcal{B}} + \alpha_1 D_x^2 \otimes id_{\mathcal{B}} + id_{\mathcal{A}} \otimes c_2 D_\xi^* \\ & + \frac{\partial H}{\partial p} D_x \otimes id_{\mathcal{B}} - id_{\mathcal{A}} \otimes (-\frac{\partial H}{\partial p_\xi}) D_\xi^*] \rho^\infty \end{aligned} \quad (41)$$

where D_ξ^* is defined by $D_\xi^* = -D_\xi L_{q^{-1}}$ ($L_{q^{-1}} f(\xi) = f(q^{-1}\xi)$ for any anyonic function $f(\xi)$) and where we have set $d_1 = d_2 = 1$. It should be pointed out though that we can express eqs.(30, 41) using the finite dimensional representation of ξ , D_ξ as we shall see in the next section.

5 Matrix realization of random walk and diffusion

We will now implement the N -dimensional matrix representations of ξ , D_ξ , D_ξ^* as they have explicitly been constructed in [23] . As to matrix representation of D_ξ^* one should take in to account the so called anyonic Leibnitz rule $D_\xi(fg) = (D_\xi f)g + L_q f(D_\xi g)$ and the property that $D_\xi^* = -D_\xi L_{q^{-1}}$ [21, 9]. Thus we have that,

$$\begin{aligned} \xi = & \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 0 & . & . & . & \dots \\ . & . & . & . & . & . & \dots \\ . & . & . & . & . & 1 & 0 \end{pmatrix}, \quad D_\xi = \begin{pmatrix} 0 & \{1\} & 0 & 0 & 0 & 0 & \dots \\ . & 0 & \{2\} & 0 & 0 & 0 & \dots \\ . & . & . & . & . & . & \dots \\ . & . & . & . & . & 0 & \{N-1\} \\ . & . & . & . & . & . & 0 \end{pmatrix}, \\ D_\xi^* = & - \begin{pmatrix} 0 & \{1\}e^{\omega^{-1}\{1\}} & 0 & 0 & 0 & 0 & \dots \\ . & 0 & \{2\}e^{\omega^{-1}\{2\}} & 0 & 0 & 0 & \dots \\ . & . & . & . & . & . & \dots \\ . & . & . & . & . & 0 & \{N-1\}e^{\omega^{-1}\{N-1\}} \\ . & . & . & . & . & . & 0 \end{pmatrix}, \end{aligned} \quad (42)$$

where $\{x\} = \frac{1-\omega^x}{1-\omega}$, $\omega = e^{2i\pi/N}$. The matrix representation of higher powers of D_ξ , D_ξ^* and ξ can easily be obtained. For a general $\rho(x, \xi)$ we have the lower triangular form

$$\begin{aligned} \rho(x, \xi, t) &= \sum_{i=0}^{N-1} \sum_{j=0}^{\infty} \rho_{ij}(t) x^j \otimes \xi^i = \sum_{i=0}^{N-1} \rho_i(x, t) \otimes \xi^i \\ &= \begin{pmatrix} \rho_0(x, t) & 0 & 0 & 0 & 0 & 0 & \dots \\ \rho_1(x, t) & \rho_0(x, t) & 0 & 0 & 0 & 0 & \dots \\ \rho_2(x, t) & \rho_1(x, t) & \rho_0(x, t) & \cdot & \cdot & \cdot & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \rho_{N-1}(x, t) & \rho_{N-2}(x, t) & \cdot & \cdot & \cdot & \rho_1(x, t) & \rho_0(x, t) \end{pmatrix}. \end{aligned} \quad (43)$$

For stationary random walks and general N -potent variables, we can now express eq.(30) in matrix form, using eqs.(42,43). The right hand side of eq.(30) becomes an operator valued upper band matrix :

$$\begin{bmatrix} H_x & c_2\lambda_1 & \alpha_2\lambda_1\lambda_2 & 0 & 0 & 0 & 0 & 0 \\ 0 & H_x & c_2\lambda_2 & \alpha_2\lambda_2\lambda_3 & 0 & 0 & 0 & 0 \\ 0 & 0 & H_x & c_2\lambda_3 & \alpha_2\lambda_3\lambda_4 & 0 & 0 & 0 \\ 0 & 0 & 0 & H_x & c_2\lambda_4 & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & H_x & \ddots & \ddots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & c_2\lambda_{N-2} & \alpha_2\lambda_{N-2}\lambda_{N-1} \\ 0 & 0 & 0 & 0 & 0 & 0 & H_x & c_2\lambda_{N-1} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & H_x \end{bmatrix}, \quad (44)$$

where $\lambda_i = \{i\}e^{\omega^{-1}\{i\}}$, $i = 1, \dots, N-1$, $H_x = -c_1D_x + \alpha_1D_x^2$, while in the left hand side $\rho^\infty(x, \xi, t)$ is as in eq.(43). The system of differential equations obtained from eqs.(43,44), describing the continue limit of stationary random walk in Ω , can be derived and is given by

$$\frac{\partial \rho_k}{\partial t} = H_x \rho_k + c_2 \lambda_{k+1} \rho_{k+1} + \alpha_2 \lambda_{k+1} \lambda_{k+2} \rho_{k+2} \quad (45)$$

for $k=0,1,\dots,N-1$. The generalization to the non stationary case is a straightforward application of the above matrix representations and the relations in eq.(41) and will not be dealt with explicitly here. Rather we shall give an explicit solution of eq.(45) for the case $N = 2$, where the above equation takes the form

$$\frac{\partial}{\partial t} \begin{bmatrix} \rho_0(x, t) \\ \rho_1(x, t) \end{bmatrix} = \begin{bmatrix} H_x & -c_2 \\ 0 & H_x \end{bmatrix} \begin{bmatrix} \rho_0 \\ \rho_1 \end{bmatrix}, \quad (46)$$

which after integration gives the time evolution

$$\begin{bmatrix} \rho_0(x, t) \\ \rho_1(x, t) \end{bmatrix} = (e^{-tH_x} + \begin{bmatrix} 1 & -tc_2 \\ 0 & 1 \end{bmatrix}) \begin{bmatrix} \rho_0(x, 0) \\ \rho_1(x, 0) \end{bmatrix}. \quad (47)$$

If initially $\rho_i(x, 0) = f_i(x) = \sum_{n \geq 0} f_{i,n} x^n$, $i = 0, 1$, then we obtain that

$$\rho'_i(x, t) = e^{-tH_x} \rho_i(x, 0) = \sum_{n \geq 0} f_{i,n} H_n(x - tc_1, t\alpha_1), \quad (48)$$

where the two variable generalized Hermite polynomial $H_n(x, y) = e^{y \frac{\partial^2}{\partial x^2}} (x^n)$ has been used. These polynomials have the generating function $e^{xt+yt^2} = \sum_{n \geq 0} \frac{t^n}{n!} H_n(x, y)$, and the expansion $H_n(x, y) = n! \sum_{r=0}^{[n/2]} \frac{y^r x^{n-2r}}{r!(n-2r)!}$. Interestingly enough these polynomials can be generated by a pair of step operators a^+ , a^- as

$$a^+ H_n(x, y) = H_{n+1}(x, y), \quad a^- H_n(x, y) = n H_{n-1}(x, y).$$

where a^+ , a^- satisfy the Heisenberg canonical commutation relation $[a^-, a^+] = 1$ and admit the realization $a^+ = x + 2xy \frac{\partial^2}{\partial x^2}$, $a^- = \frac{\partial}{\partial x}$ (see e.g. [5]).

Finally the solution of eq.(46) reads

$$\begin{bmatrix} \rho_0(x, t) \\ \rho_1(x, t) \end{bmatrix} = \begin{bmatrix} \rho'_0(x, t) + \rho_0(x, 0) - tc_2 \rho_1(x, 0) \\ \rho'_1(x, t) + \rho_1(x, 0) \end{bmatrix}.$$

6 Transition operators

In this chapter we shall formulate Markov processes on the smash line algebra. The key point is to define the Markov transition operator $T_\phi : \Omega \rightarrow \Omega$, where ϕ is as before a linear functional on Ω , such that

$$T_\phi = (\phi \otimes id_\Omega)\Delta_\Omega, \quad \varepsilon_\Omega \circ T_\phi = \phi, \quad \phi * \psi = \varepsilon_\Omega \circ T_\psi T_\phi. \quad (49)$$

Last relation implies that the Markov transition operators T_ϕ , can be composed to form a semigroup with $T_{\phi=\varepsilon_\Omega} = id_\Omega$, as unit element, which is homomorphic to the convolution semigroup of functionals with unit element the counit map ε_Ω , and generator the functional ϕ . In this active picture of random walk in terms of transition operators, to find the expectation values of the observables after n steps we employ the relation

$$\langle f \rangle_{\phi_1 * \phi_2 * \dots * \phi_n} = \varepsilon_\Omega \circ T_{\phi_1} T_{\phi_2} \dots T_{\phi_n}(f),$$

where in general the states ϕ are varying at each step. Let $f(x, \xi)$ be any function smooth of x and ξ . We shall look at stationary random walks first. Using eq.(49) with $\phi(f) = \int \rho f$, and probability density as $\rho(x, \xi) = \rho_1(x) \otimes \rho_2(\xi)$, with $\rho_1(x)$, $\rho_2(\xi)$ as in eq.(18), we obtain that after one step

$$\begin{aligned} (T_\phi f)(x, \xi) &= p_1 p_2 f(x + a, \xi + \theta) + (1 - p_1) p_2 f(x - a, \xi + \theta) \\ &\quad + p_1 (1 - p_2) f(x + a, \xi - \theta) + (1 - p_1) (1 - p_2) f(x - a, \xi - \theta) \\ &= [p_1 e^{aD_x} \otimes id_{\mathcal{B}} + (1 - p_1) e^{-aD_x} \otimes id_{\mathcal{B}}] \\ &\quad \times [p_2 id_{\mathcal{A}} \otimes e^{\theta D_\xi} + (1 - p_2) id_{\mathcal{A}} \otimes e^{-\theta D_\xi}] f(x, \xi) \end{aligned} \quad (50)$$

$$= (T_{\phi_x} \otimes T_{\phi_\xi}) f(x, \xi), \quad (51)$$

while when $\rho(x, \xi) = \frac{1}{2}[\rho_1(x) \otimes I_{\mathcal{B}} + I_{\mathcal{A}} \otimes \rho_2(\xi)]$, we obtain

$$\begin{aligned} (T_\phi f)(x, \xi) &= p_1 f(x + a, \xi) + (1 - p_1) f(x - a, \xi) + \\ &\quad p_2 f(x, \xi + \theta) + (1 - p_2) f(x, \xi - \theta) \\ &= [p_1 e^{aD_x} \otimes id_{\mathcal{B}} + (1 - p_1) e^{-aD_x} \otimes id_{\mathcal{B}}] \end{aligned} \quad (52)$$

$$\begin{aligned} &\quad + [p_2 id_{\mathcal{A}} \otimes e^{\theta D_\xi} + (1 - p_2) id_{\mathcal{A}} \otimes e^{-\theta D_\xi}] f(x, \xi) \\ &= (T_{\phi_x} \otimes id_{\mathcal{B}} + id_{\mathcal{A}} \otimes T_{\phi_\xi}) f(x, \xi). \end{aligned} \quad (53)$$

The Markov transition operator in the continue limit T_{ϕ^∞} can be obtained from eq.(50) with exactly the same procedure of Taylor expansion and the use of substitutions as in eq.(24), this yields the operator

$$T_{\phi^\infty} = e^{(-c_1 t D_x + \alpha_1 t D_x^2) \otimes id_{\mathcal{B}} + id_{\mathcal{A}} \otimes (c_2 t D_\xi + \alpha_2 t D_\xi^2)}. \quad (54)$$

We can also defined T_ϕ in the non stationary case where ϕ depends on time (or equivalently, as we have already seen, when x, ξ themselves evolve in time) and a straightforward calculation using eq.(37,49), show that for the case of the real line merged with Grassmann line i.e $N = 2$,

$$T_{\phi_t} = (\phi_t \otimes id_\Omega) \circ \Delta_\Omega = [e^{-\lambda t \partial / \partial a} \otimes (1 - \tilde{\lambda} t \partial / \partial \theta)] T_{\phi_{t=0}}, \quad (55)$$

where $T_{\phi_{t=0}}$ is as in eq.(50). Taking the continue limit of the non stationary case, as we did in the case of ϕ , leads to

$$T_{\phi_t}^\infty = e^{(-c_1 + \lambda) t D_x \otimes id_{\mathcal{B}} + \alpha_1 t D_x^2 \otimes id_{\mathcal{B}} + id_{\mathcal{A}} \otimes (c_2 - \tilde{\lambda}) t D_\xi^*}. \quad (56)$$

Concluding this section we must point out that we can have a left and a right transition operators T_ϕ^L, T_ϕ^R , defined respectively by eq.(49) and $T_\phi^R = (id_\Omega \otimes \phi) \circ \Delta_\Omega$, and so the above constructions have a left and right version. Also in the case where the functional is identified with the counit i.e $\phi = \varepsilon_\Omega = \int$, the respective transition operators $T_{\phi=\varepsilon_\Omega}^L = T_{\phi=\varepsilon_\Omega}^R = id_\Omega$. and their dual probability density $\rho_{\phi=\varepsilon_\Omega} = I_\Omega$, is trivial.

7 Entanglement and statistical correlations

We are now in a position to generalize the procedure of the previous sections to obtain diffusion equations when the choice of the density $\rho(x, \xi)$ has the general form of a convex combination as:

$$\rho(x, \xi) = \sum_{i=1}^m \lambda_i \rho_1^i(x) \otimes \rho_2^i(\xi), \quad \lambda_i \geq 0, \quad \sum_{i=1}^m \lambda_i = 1. \quad (57)$$

From relations (13) and (15) we have that

$$\phi_\Omega = \sum_{i=1}^m \lambda_i \phi_x^i \otimes \phi_\xi^i, \quad (58)$$

which in the case $m = 1$ and thus $\lambda_1 = 1$, $\rho(x, \xi)$ reduces to the form of eq.(23) which has explicitly dealt with in the previous chapters. The choice $m = 2$ and thus $\lambda_1 + \lambda_2 = 1$, $\rho(x, \xi)$ reduces to the form

$$\rho_\Omega(x, \xi) = \lambda \rho_1(x) \otimes I_{\mathcal{B}} + (1 - \lambda) I_{\mathcal{A}} \otimes \rho_2(\xi) \quad (59)$$

which implies that

$$\phi_\Omega = \lambda \phi_x \otimes \varepsilon_{\mathcal{B}} + (1 - \lambda) \varepsilon_{\mathcal{A}} \otimes \phi_\xi \quad (60)$$

Making the choices $\rho_1^i(x) = p_1^i \delta(x - a^i) + (1 - p_1^i) \delta(x + a^i)$, $\rho_2^i(\xi) = p_2^i \delta(\xi - \theta^i) + (1 - p_2^i) \delta(\xi + \theta^i)$, the general choice of eq.(57) leads to the following Taylor expanded form of $\phi_\Omega^{\star n}$:

$$\begin{aligned} \phi_\Omega^{\star n} &= \left[\sum_{i=1}^m \lambda_i (p_1^i e^{a^i D_x} + (1 - p_1^i) e^{-a^i D_x}) \otimes (p_2^i e^{\theta^i D_\xi} + (1 - p_2^i) e^{-\theta^i D_\xi}) \right]_{|x=0, \xi=0}^n \\ &= \left[\sum_{i=1}^m \lambda_i (\varepsilon_\Omega + 2a^i (p_1^i - 1/2) D_x \otimes \varepsilon_{\mathcal{B}} + a^{i2}/2! D_x^2 \otimes \varepsilon_{\mathcal{B}} + \dots) \right. \\ &\quad \left. \star (\varepsilon_\Omega + \varepsilon_{\mathcal{A}} \otimes 2\theta^i (p_2^i - 1/2) D_\xi + \varepsilon_{\mathcal{A}} \otimes (\theta^i)^2/[2]_q! D_\xi^2 + \dots) \right]_{|x=0, \xi=0}^n \end{aligned}$$

We now make the substitutions

$$\begin{aligned} 2a^i (p_1^i - 1/2) &= \frac{c_1^i t}{n}, \quad 2\theta^i (p_2^i - 1/2) = \frac{c_2^i t}{n}, \\ (a^i)^2/2 &= \frac{\alpha_1^i t}{n}, \quad (\theta^i)^2/[2]_q = \frac{\alpha_2^i t}{n}, \end{aligned} \quad (61)$$

which lead to

$$\begin{aligned}\phi_{\Omega}^{\star n} = & \left\{ \sum_{i=1}^m \lambda_i [\varepsilon_{\Omega} + \frac{1}{n} (c_1^i t D_x + \alpha_1^i t D_x^2) \otimes \varepsilon_{\mathcal{B}} + \frac{1}{n} \varepsilon_{\mathcal{A}} \otimes (c_2^i t D_{\xi} + \alpha_2^i t D_{\xi}^2) \right. \\ & \left. + \frac{t^2}{n^2} (c_1^i D_x + \alpha_1^i D_x^2) (c_2^i D_{\xi} + \alpha_2^i D_{\xi}^2) \right\}_{|x=0, \xi=0}^n.\end{aligned}$$

By implementing the limit $\lim_{n \rightarrow \infty} (1 + z/n + w/n^2) = e^z$, it is easily obtained that

$$\phi_{\Omega}^{\infty} = \exp t \left\{ \sum_{i=1}^m \lambda_i [(c_1^i D_x + \alpha_1^i D_x^2) \otimes \varepsilon_{\mathcal{B}} + \varepsilon_{\mathcal{A}} \otimes (c_2^i D_{\xi} + \alpha_2^i D_{\xi}^2)] \right\}_{|x=0, \xi=0}. \quad (62)$$

Defining the drift and diffusion terms as $L_{\mathcal{A},i}^{drift} = c_1^i D_x \otimes \varepsilon_{\mathcal{B}}$, $L_{\mathcal{B},i}^{drift} = \varepsilon_{\mathcal{A}} \otimes c_2^i D_{\xi}$, $L_{\mathcal{A},i}^{diff} = \alpha_1^i D_x^2 \otimes \varepsilon_{\mathcal{B}}$, and $L_{\mathcal{B},i}^{diff} = \varepsilon_{\mathcal{A}} \otimes \alpha_2^i D_{\xi}^2$, eq.(62) can be written as

$$\phi_{\Omega}^{\infty} = \exp t \left\{ \sum_{i=1}^m \lambda_i (L_{\mathcal{A},i}^{drift} + L_{\mathcal{B},i}^{drift} + L_{\mathcal{A},i}^{diff} + L_{\mathcal{B},i}^{diff}) \right\}_{|x=0, \xi=0}. \quad (63)$$

The form of the transition operator in this general case is shown to be

$$T_{\Omega}^{\infty} = \exp t \left\{ \sum_{i=1}^m \lambda_i [(-c_1^i D_x + \alpha_1^i D_x^2) \otimes id_{\mathcal{B}} + id_{\mathcal{A}} \otimes (c_2^i D_{\xi}^* + \alpha_2^i D_{\xi}^{*2})] \right\}, \quad (64)$$

while the diffusion equation reads

$$\frac{\partial \rho_{\Omega}^{\infty}}{\partial t} = \sum_{i=1}^m \lambda_i (-L_{\mathcal{A},i}^{drift} + L_{\mathcal{B},i}^{drift} + L_{\mathcal{A},i}^{diff} + L_{\mathcal{B},i}^{*diff}) \rho_{\Omega}^{\infty}, \quad (65)$$

where $L_{\mathcal{A},i}^{drift} = c_1^i D_x \otimes id_{\mathcal{B}}$, $L_{\mathcal{B},i}^{*drift} = id_{\mathcal{A}} \otimes c_2^i D_{\xi}^*$, $L_{\mathcal{A},i}^{diff} = \alpha_1^i D_x^2 \otimes id_{\mathcal{B}}$, and $L_{\mathcal{B},i}^{*diff} = id_{\mathcal{A}} \otimes \alpha_2^i D_{\xi}^{*2}$. The above equation can be rewritten as

$$\frac{\partial \rho_{\Omega}^{\infty}}{\partial t} = (-c_1 D_x \otimes id_{\mathcal{B}} + \alpha_1 D_x^2 \otimes id_{\mathcal{B}} + c_2 id_{\mathcal{A}} \otimes D_{\xi}^* + \alpha_2 id_{\mathcal{A}} \otimes D_{\xi}^{*2}) \rho_{\Omega}^{\infty} \quad (66)$$

where

$$c_1 = \sum_{i=1}^m \lambda_i c_1^i, \quad c_2 = \sum_{i=1}^m \lambda_i c_2^i, \quad \alpha_1 = \sum_{i=1}^m \lambda_i \alpha_1^i, \quad \alpha_2 = \sum_{i=1}^m \lambda_i \alpha_2^i. \quad (67)$$

For uncorrelated walks the density function is factorized i.e $\rho(x, \xi) = \rho_1(x) \otimes \rho_2(\xi)$, and in that case we obtain the statistical moments

$$\begin{aligned} \mu_{kl} &= \langle x^k \otimes \xi^l \rangle_{\phi_\Omega} = \int \rho(x, \xi) x^k \otimes \xi^l \\ &= \int \rho_1(x) x^k \otimes \rho_2(\xi) \xi^l = \phi_x(x^k) \phi_\xi(\xi^l) = \mu_k^x \mu_l^\xi, \end{aligned} \quad (68)$$

where $0 \leq k < \infty$, $0 \leq l \leq N-1$ and $\mu_k^x = \phi_x(x^k)$, $\mu_l^\xi = \phi_\xi(\xi^l)$. For density functions of the form $\rho(x, \xi) = \lambda \rho_1(x) \otimes I_{\mathcal{B}} + (1-\lambda) I_{\mathcal{A}} \otimes \rho_2(\xi)$ we have that

$$\begin{aligned} \mu_{kl} &= \langle x^k \otimes \xi^l \rangle_{\phi_\Omega} = \int \rho(x, \xi) x^k \otimes \xi^l = \lambda \phi_x(x^k) + (1-\lambda) \phi_\xi(\xi^l) \\ &= \lambda \mu_k^x + (1-\lambda) \mu_l^\xi. \end{aligned} \quad (69)$$

Finally, for classically correlated or seperable density functions

$$\rho(x, \xi) = \sum_{i=1}^m \lambda_i \rho_i(x) \otimes \rho_i(\xi), \quad \lambda_i \geq 0, \quad \sum_{i=1}^m \lambda_i = 1, \quad (70)$$

we get

$$\mu_{kl} = \sum_{i=1}^m \lambda_i \phi_x^i(x^k) \phi_\xi^i(\xi^l) = \sum_{i=1}^m \lambda_i \mu_k^{x_i} \mu_l^{\xi_i}. \quad (71)$$

We can now proceed to obtain a differential equation for the moments. By evaluating $\frac{\partial}{\partial t} \langle x^k \rangle_{\phi_\Omega^\infty}$, $\frac{\partial}{\partial t} \langle \xi^l \rangle_{\phi_\Omega^\infty}$ and $\frac{\partial}{\partial t} \langle x^k \otimes \xi^l \rangle_{\phi_\Omega^\infty}$, we obtain the equations

$$\begin{aligned}
\frac{\partial}{\partial t} \mu_{k,l}(t) &= \frac{\partial}{\partial t} \langle x^k \otimes \xi^l \rangle_{\phi_{\Omega}^{\infty}} = \int (x^k \otimes \xi^l) \left(\frac{\partial}{\partial t} \rho_{\Omega}^{\infty} \right) \\
&= \sum_{i=1}^m \lambda_i \left[\int (x^k \otimes \xi^l) (-L_{\mathcal{A},i}^{drift} + L_{\mathcal{B},i}^{drift} + L_{\mathcal{A},i}^{*diff} + L_{\mathcal{B},i}^{*diff}) \rho_{\Omega}^{\infty} \right] \\
&= \sum_{i=1}^m \lambda_i [c_1^i k \mu_{k-1,l}(t) + \alpha_1^i k(k-1) \mu_{k-2,l}(t) \\
&\quad + c_2^i [l]_q \mu_{k,l-1}(t) + \alpha_2^i [l]_q [l-1]_q \mu_{k,l-2}(t)] \\
&= c_1 k \mu_{k-1,l}(t) + \alpha_1 k(k-1) \mu_{k-2,l}(t) + c_2 [l]_q \mu_{k,l-1}(t) + \alpha_2 [l]_q [l-1]_q \mu_{k,l-2}(t),
\end{aligned} \tag{72}$$

where in the above derivation integration by parts has also been used for the q-Jackson integral involving ξ s [10], and $c_1, \alpha_1, c_2, \alpha_2$, are given in eq.(67).

In particular this last formula can be cast in two interesting reduced forms when we consider only x -type diffusion or the ξ -type one. For the first case, if we introduce the column vector of x -statistical moments $\kappa \equiv (\mu_k)_{k \geq 0}$ where $\mu_k \equiv \mu_{k,0}$, and invoke the matrix representation of the canonical Heisenberg algebra generated by $a, a^\dagger$ and I , where the number operator is $N = a^\dagger a$, we can express the dynamics of the moments by means of the relation

$$\frac{\partial}{\partial t} \kappa(t) = (c_1 a \sqrt{N} + \alpha_1 a \sqrt{N} a \sqrt{N}) \kappa. \tag{73}$$

Similarly for the ξ -type case we have that

$$\frac{\partial}{\partial t} \nu = (c_2 a_q \sqrt{[N_q]_q} + \alpha_2 a_q \sqrt{[N_q]_q} a_q \sqrt{[N_q]_q}) \nu, \tag{74}$$

where c_1, α_1 , and c_2, α_2 , are given in eq.(67), and the column vector of the ξ -statistical moments $\nu \equiv (\nu)_{l=0}^{N-1} = (\mu_{0,l})_{l=0}^{N-1}$, is introduced. The following operators

$$a_q = \begin{pmatrix} 0 & \sqrt{[1]_q} & 0 & 0 & 0 & 0 & \dots \\ \cdot & 0 & \sqrt{[2]_q} & 0 & 0 & 0 & \dots \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \dots \\ \cdot & \cdot & \cdot & \cdot & \cdot & 0 & \sqrt{[N-1]_q} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 0 \end{pmatrix}, \quad N_q = - \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ \cdot & 1 & 0 & 0 & 0 & 0 & \dots \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \dots \\ \cdot & \cdot & \cdot & \cdot & \cdot & N-2 & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & N-1 \end{pmatrix},$$

together with the conjugate $a_q^\dagger$ of a_q , form a truncated N -dimensional matrix representation of the so called q -canonical algebra for $q = e^{i2\pi/N}$ [7].

Closing we point out that eq. (72) can also be cast in the following form

$$\frac{\partial}{\partial t}\mu = (\mathbf{1}_N \otimes L_x + L_\xi \otimes \mathbf{1}_\infty)\mu$$

where $\mathbf{1}_N$ is a $N \times N$ unit matrix, $\mathbf{1}_\infty$ is an infinite dimensional unit matrix, $L_x = c_1 a \sqrt{N} + \alpha_1 a \sqrt{N} a \sqrt{N}$, and $L_\xi = c_2 a_q \sqrt{[N_q]_q} + \alpha_2 a_q \sqrt{[N_q]_q} a_q \sqrt{[N_q]_q}$. Also μ is a column vector given in terms of the statistical moments of the two partial random walks viz.

$$\mu = [\mu_{00}, \mu_{10}, \mu_{20}, \dots | \mu_{01}, \mu_{11}, \mu_{21}, \dots | \dots | \mu_{0N-1}, \mu_{1N-1}, \mu_{2N-1}, \dots]^T.$$

8 Conclusion

In this paper we have used the machinery of Hopf algebras and the algebraic braiding or smashing between algebras, in order to construct a non trivial extension of a 1D random walk. In the continuous limit its diffusion equation has been constructed and a matrix representation of it has been obtained and solved for some spacial cases. Emphasis has been put on three aspects: firstly, the role of algebra braiding in connection with the type of non commutativity among the steps of the random walk, secondly, the role of entanglement (statistical correlations) among the density probability functions of the two partial random walks with variables x and ξ and thirdly, the role of canonical and deformed Heisenberg algebra in the equations ruling the temporal evolution of the statistical moments of the 2D random walk.

Acknowledgments

One of us (I. T.) would like to thank the National Foundation of Scholarships of Greece (IKY) for financial support.

References

- [1] E. Abe, *Hopf Algebras* (CUP Cambridge 1997).
- [2] L. Accardi, M. Schürman and W. von Waldenfels, Math. Z. **198**, 451-477 (1988).
- [3] Y. Bessalov and B. Drabant, *Cross Product Bialgebras - Part I-II*, math/9802028, math/9904142.
- [4] S. Caenepeel, I. Bogdan, G. Militaru and S. Zhu, *Smash biproducts of algebras and coalgebras*, math.qa-9809063;
- [5] G. Dattoli, J. Computational and Applied Math. **118**, 111 (2000).
- [6] E. D. Davies, *Quantum Theory of Open System*, (Academic, New York, 1973).
- [7] D. Ellinas, Phys. Rev. A **45**, 3358-3361 (1992).
- [8] D. Ellinas, J. Comp. and Applied Math. **133** 341-353 (2001).
- [9] D. Ellinas and I. Tsochantjis, J. of Non-Linear Math. Physics Supp. **8**, 100 (2001).
- [10] H. Exton, *q-Hypergeometric Functions and Applications*, (Horwood, Chichester 1983).
- [11] P. Feinsilver and R. Schott, J. Theor. Prob. **5**, 251 (1992).
- [12] P. Feinsilver, U. Franz and R. Schott, J. Theor. Prob. **10**, 797 (1997).
- [13] U. Franz and R. Schott, *Stochastic Processes and Operator Calculus on Quantum Groups*, (Kluwer Academic Publishers, Dordrecht, 1999).
- [14] U. Franz and R. Schott, J. Phys. A: Gen. Math. **31**, 1395 (1998).
- [15] U. Franz and R. Schott, J. Math. Phys. **39**, 2748 (1998).

- [16] N. Giri and W. von Waldenfels, *Z. Wahr. Verw. Gebiete* **42**, 129-134 (1978).
- [17] A. P. Isaev, *Paragrassmann Integral, Discrete Systems and Quantum Groups*, q-alg/9609030.
- [18] R. Lenczewski, *Comm. Math. Phys.* **154**, 127-134 (1993).
- [19] S. Majid, *Foundations of Quantum Groups Theory* (Cambridge Univ. Press, 1995), ff. chapter 5.
- [20] S. Majid, *Int. J. Mod. Phys.* **8**, 4521-4545 (1993).
- [21] S. Majid and M. J. Rodriguez-Plaza, *J. Math. Phys.* **33**, 3753-3760 (1994).
- [22] P. A. Meyer, *Quantum Probability for Probabilists* (Lect. Notes Math. 1538), (Springer, Berlin 1993).
- [23] M. Rausch de Traubenberg, *Adv. Appl. Clifford Alg.* **4**, 131 (1994).
- [24] M. Schürman, *White Noise on Bialgebras* (Lect. Notes Math. 1544), (Springer, Berlin 1993).
- [25] M. Schürman, *Lect. Notes Math.* **1210**, 153-157 (1986).
- [26] M. Schürman, *Comm. Math. Phys.* **140**, 589-615 (1991).
- [27] R. Speicher, *Prob. Th. Rel. Fields* **84**, 141-159 (1990).
- [28] M. E. Sweedler, *Hopf Algebras*, (Benjamin, New York 1969).
- [29] D. Voiculescu, *J. Funct. Anal.* **66**, 323-346 (1986).
- [30] W. von Waldenfels, *Z. Wahr. Verw. Gebiete* **42**, 135-140 (1978).